

# Staying Healthy: Recommendations for MEN



The Veterans Health Administration (VHA) aims to help you stay healthy. The table below lists the **preventive health services** (screening tests, medications, health counseling, and vaccines) that VHA recommends. Screening tests are used to look for health conditions before there are symptoms. These recommendations apply only to adult men of **average risk**. You are a man of average risk if you have **no personal or family history or symptoms** of the conditions listed below. If you are having symptoms of a condition, please talk with your provider.

## Which preventive health services are recommended for average risk men in my age group?

Medium gray:	Recommended
Light gray:	Recommended for some men – talk with your provider
Dark gray:	Not recommended (NR)

## SCREENING TESTS FOR AVERAGE RISK VETERANS

HEALTH CONDITIONS	18–29 years	30–39 years	40–49 years	50–59 years	60–69 years	70–79 years	80 years and older	
Abdominal Aortic Aneurysm (AAA)	Not recommended (age 18–64)					Talk with your provider (age 65–75)	Not recommended (age 76 and older)	
Colon Cancer	Not recommended (age 18–44)			Recommended. Frequency varies by test chosen (age 45–75)			Talk with your provider (age 76–85)	NR (age 86 and older)
Depression	Recommended every year							
Hepatitis B Infection	Recommended for some men – talk with your provider (age 18 and older)							
Hepatitis C Infection	Recommended in adults aged 18-79 years							
High Blood Pressure	Recommended every year (age 18 and older)							
High Cholesterol		Talk with your provider (age 20 and older)						
HIV Infection	Recommended once (age 18 and older); annually in Veterans with ongoing risk factors							
Lung Cancer	Not recommended (age 18–49)			Recommended for some men – talk with your provider (age 50–80)				Not recommended (age 81 and older)
Overweight & Obesity	Recommended every year (age 18 and older)							
Prostate Cancer	Not recommended (age 18–54)				Talk with your provider (age 55-69)		Not recommended (age 70 and older)	
Sexually Transmitted Infections	Talk with your provider about testing for syphilis and other sexually transmitted infections (age 18 and older)							

## MEDICATIONS

HEALTH CONDITIONS	18–29 years	30–39 years	40–49 years	50–59 years	60–69 years	70–79 years	80 years and older
Aspirin to Prevent Heart Attack	Not recommended (age 18–49)			Talk with your provider (age 50–69)		Not recommended (age 70 and older)	

## HEALTH COUNSELING

HEALTH CONDITIONS	18–29 years	30–39 years	40–49 years	50–59 years	60–69 years	70–79 years	80 years and older
Tobacco Use	Recommended every visit (if using tobacco) (age 18 and older)						
Alcohol Use	Talk with your provider about healthy alcohol use (age 18 and older)						
Healthy Diet and Physical Activity	Talk with your provider about a healthy diet and physical activity (age 18 and older)						

## VACCINES

HEALTH CONDITIONS	18–29 years	30–39 years	40–49 years	50–59 years	60–69 years	70–79 years	80 years and older
Hepatitis A	Recommended for some men – talk with your provider (age 18 and older)						
Hepatitis B	Recommended for some men – talk with your provider (age 18 and older)						
Herpes Zoster (Shingles) RZV vaccine	Not recommended (age 18–49)			Recommended 2 doses (age 50 and older), with second dose 2–6 months after first dose			
Human Papilloma Virus (HPV) HPV9 vaccine	2–3 doses (age 18–26) if series not completed	Talk with your provider (age 27–45)		Not recommended (age 46 and older)			
Influenza (Flu) Injectable, Inhaled	Recommended every year (age 18 and older)						
Measles, Mumps, and Rubella (MMR)	Recommended for some men – talk with your provider (age 18 and older)						
Meningococcal Disease	Recommended for some men – talk with your provider (age 18 and older)						
Pneumococcal Disease PPSV23 vaccine, PCV13 vaccine	Recommended for some men – talk with your provider (age 18–64)				Recommended. At least 1 dose of PPSV23 vaccine (age 65 and older)		
					Talk with your provider about PCV13 vaccine (age 65 and older)		
Tdap (Tetanus, Diphtheria & Pertussis)	Recommend at least 1 dose (age 18 and older) (Or for wound management if > 5 years have passed since last tetanus toxoid-containing booster vaccine).						
Td (Tetanus and diphtheria)	Recommended. Booster every 10 years (age 18 and older) (Either Tdap or Td may be used) (Or for wound management if > 5 years have passed since last tetanus toxoid-containing booster vaccine).						
Varicella (Chickenpox)	Recommended for some men - talk with your provider (2 doses if born 1980 or later)						

